

Таблица связи обратных тригонометрических функций

	<i>arcsin y</i>	<i>arccos y</i>	<i>arctg y</i>	<i>arcctg y</i>
<i>arcsin x</i>		$y = \sqrt{1-x^2}$ $0 \leq x \leq 1$	$y = \frac{x}{\sqrt{1-x^2}}$ $0 < x < 1$	$y = \frac{\sqrt{1-x^2}}{x}$ $0 < x \leq 1$
<i>arccos x</i>	$y = \sqrt{1-x^2}$ $0 \leq x \leq 1$		$y = \frac{\sqrt{1-x^2}}{x}$ $0 < x \leq 1$	$y = \frac{x}{\sqrt{1-x^2}}$ $0 < x < 1$
<i>arctg x</i>	$y = \frac{x}{\sqrt{1+x^2}}$ $x \in \mathbb{R}$	$y = \frac{1}{\sqrt{1+x^2}}$ $x \geq 0$		$y = \frac{1}{x}$ $x > 0$
<i>arccg x</i>	$y = \frac{1}{\sqrt{1+x^2}}$ $x \geq 0$	$y = \frac{x}{\sqrt{1+x^2}}$ $x \in \mathbb{R}$	$y = \frac{1}{x}$ $x > 0$	

I. Арксинус

1. $\arcsin x = \arccos y = t;$

$$\begin{cases} -\frac{\pi}{2} \leq \arcsin x \leq \frac{\pi}{2}, \\ 0 \leq \arccos y \leq \pi; \end{cases} \Rightarrow 0 \leq t \leq \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 \leq \arcsin x \leq \frac{\pi}{2}, \\ 0 \leq \arccos y \leq \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} 0 \leq x \leq 1, \\ 0 \leq y \leq 1. \end{cases}$$

$$\begin{cases} \arcsin x = t, \\ \arccos y = t; \end{cases} \Leftrightarrow \begin{cases} x = \sin t, \\ y = \cos t. \end{cases}; \quad 0 \leq t \leq \frac{\pi}{2} \Rightarrow \cos t = \sqrt{1 - \sin^2 t} \Rightarrow y = \sqrt{1 - x^2}.$$

2. $\arcsin x = \operatorname{arctg} y = t;$

$$\begin{cases} -\frac{\pi}{2} \leq \arcsin x \leq \frac{\pi}{2}, \\ -\frac{\pi}{2} < \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Rightarrow -\frac{\pi}{2} < t < \frac{\pi}{2} \Leftrightarrow \begin{cases} -\frac{\pi}{2} < \arcsin x < \frac{\pi}{2}, \\ -\frac{\pi}{2} < \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} 0 < x < 1, \\ y \in \mathbb{R}. \end{cases}$$

$$\begin{cases} \arcsin x = t, \\ \operatorname{arctg} y = t; \end{cases} \Leftrightarrow \begin{cases} x = \sin t, \\ y = \operatorname{tg} t. \end{cases}; \quad -\frac{\pi}{2} < t < \frac{\pi}{2} \Rightarrow \operatorname{tg} t = \frac{\sin t}{\sqrt{1 - \sin^2 t}} \Rightarrow y = \frac{x}{\sqrt{1 - x^2}}.$$

3. $\arcsin x = \operatorname{arctg} y = t;$

$$\begin{cases} -\frac{\pi}{2} \leq \arcsin x \leq \frac{\pi}{2}, \\ 0 < \operatorname{arctg} y < \pi; \end{cases} \Rightarrow 0 < t \leq \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 < \arcsin x \leq \frac{\pi}{2}, \\ 0 < \operatorname{arctg} y \leq \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} 0 < x \leq 1, \\ y \geq 0. \end{cases}$$

$$\begin{cases} \arcsin x = t, \\ \operatorname{arctg} y = t; \end{cases} \Leftrightarrow \begin{cases} x = \sin t, \\ y = \operatorname{ctg} t. \end{cases}; \quad 0 < t \leq \frac{\pi}{2} \Rightarrow \operatorname{ctg} t = \frac{\sqrt{1 - \sin^2 t}}{\sin t} \Rightarrow y = \frac{\sqrt{1 - x^2}}{x}.$$

II. Арккосинус

1. $\arccos x = \arcsin y = t;$

$$\begin{cases} 0 \leq \arccos x \leq \pi, \\ -\frac{\pi}{2} \leq \arcsin y \leq \frac{\pi}{2}; \end{cases} \Rightarrow 0 \leq t \leq \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 \leq \arccos x \leq \frac{\pi}{2}, \\ 0 \leq \arcsin y \leq \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} 0 \leq x \leq 1, \\ 0 \leq y \leq 1. \end{cases}$$

$$\begin{cases} \arccos x = t, \\ \arcsin y = t; \end{cases} \Leftrightarrow \begin{cases} x = \cos t, \\ y = \sin t. \end{cases}; \quad 0 \leq t \leq \frac{\pi}{2} \Rightarrow \sin t = \sqrt{1 - \cos^2 t} \Rightarrow y = \sqrt{1 - x^2}.$$

2. $\arccos x = \operatorname{arctg} y = t;$

$$\begin{cases} 0 \leq \arccos x \leq \pi, \\ -\frac{\pi}{2} < \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Rightarrow 0 \leq t < \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 \leq \arccos x < \frac{\pi}{2}, \\ 0 \leq \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} 0 < x \leq 1, \\ y \geq 0. \end{cases};$$

$$\begin{cases} \arccos x = t, \\ \operatorname{arctg} y = t; \end{cases} \Leftrightarrow \begin{cases} x = \cos t, \\ y = \operatorname{tg} t. \end{cases}; \quad 0 \leq t < \frac{\pi}{2} \Rightarrow \operatorname{tg} t = \frac{\sqrt{1 - \cos^2 t}}{\cos t} \Rightarrow y = \frac{\sqrt{1 - x^2}}{x}.$$

3. $\arccos x = \operatorname{arctg} y = t;$

$$\begin{cases} 0 \leq \arccos x \leq \pi, \\ 0 < \operatorname{arctg} y < \pi; \end{cases} \Rightarrow 0 < t < \pi \Leftrightarrow \begin{cases} 0 < \arccos x < \pi, \\ 0 < \operatorname{arctg} y < \pi; \end{cases} \Leftrightarrow \begin{cases} -1 < x < 1, \\ y \in \mathbb{R}. \end{cases};$$

$$\begin{cases} \arccos x = t, \\ \operatorname{arctg} y = t; \end{cases} \Leftrightarrow \begin{cases} x = \cos t, \\ y = \operatorname{tg} t. \end{cases}; \quad 0 < t < \pi \Rightarrow \operatorname{tg} t = \frac{\sin t}{\cos t} \Rightarrow y = \frac{x}{\sqrt{1 - x^2}}.$$

III. Арктангенс

1. $\operatorname{arctg} x = \arcsin y = t;$

$$\begin{cases} -\frac{\pi}{2} < \operatorname{arctg} x < \frac{\pi}{2}, \\ -\frac{\pi}{2} \leq \arcsin y \leq \frac{\pi}{2}; \end{cases} \Rightarrow -\frac{\pi}{2} < t < \frac{\pi}{2} \Leftrightarrow \begin{cases} -\frac{\pi}{2} < \operatorname{arctg} x < \frac{\pi}{2}, \\ -\frac{\pi}{2} < \arcsin y < \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} x \in \mathbb{R}, \\ -1 < y < 1. \end{cases};$$

$$\begin{cases} \operatorname{arctg} x = t, \\ \arcsin y = t; \end{cases} \Leftrightarrow \begin{cases} x = \operatorname{tg} t, \\ y = \sin t. \end{cases}; \quad -\frac{\pi}{2} < t < \frac{\pi}{2} \Rightarrow \sin t = \frac{\operatorname{tg} t}{\sqrt{1 + \operatorname{tg}^2 t}} \Rightarrow y = \frac{x}{\sqrt{1 + x^2}}.$$

2. $\operatorname{arctg} x = \arccos y = t;$

$$\begin{cases} -\frac{\pi}{2} < \operatorname{arctg} x < \frac{\pi}{2}, \\ 0 \leq \arccos y \leq \pi; \end{cases} \Rightarrow 0 \leq t < \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 \leq \operatorname{arctg} x < \frac{\pi}{2}, \\ 0 \leq \arccos y < \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} x \geq 0, \\ 0 < y \leq 1. \end{cases};$$

$$\begin{cases} \operatorname{arctg} x = t, \\ \arccos y = t; \end{cases} \Leftrightarrow \begin{cases} x = \operatorname{tg} t, \\ y = \cos t. \end{cases}; \quad 0 \leq t < \frac{\pi}{2} \Rightarrow \cos t = \frac{1}{\sqrt{1 + \operatorname{tg}^2 t}} \Rightarrow y = \frac{1}{\sqrt{1 + x^2}}.$$

$$3. \operatorname{arctg} x = \operatorname{arctg} y = t;$$

$$\begin{cases} -\frac{\pi}{2} < \operatorname{arctg} x < \frac{\pi}{2}, \\ 0 < \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Rightarrow 0 < t < \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 < \operatorname{arctg} x < \frac{\pi}{2}, \\ 0 < \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} x > 0, \\ y > 0. \end{cases};$$

$$\begin{cases} \operatorname{arctg} x = t, \\ \operatorname{arctg} y = t; \end{cases} \Leftrightarrow \begin{cases} x = \operatorname{tg} t, \\ y = \operatorname{tg} t. \end{cases}; \operatorname{tg} t = \frac{1}{\operatorname{ctg} t} \Rightarrow y = \frac{1}{x}.$$

IV. Арккотангенс

$$1. \operatorname{arctg} x = \arcsin y = t;$$

$$\begin{cases} 0 < \operatorname{arctg} x < \frac{\pi}{2}, \\ -\frac{\pi}{2} \leq \arcsin y \leq \frac{\pi}{2}; \end{cases} \Rightarrow 0 < t \leq \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 < \operatorname{arctg} x \leq \frac{\pi}{2}, \\ 0 < \arcsin y \leq \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} x \geq 0, \\ 0 < y \leq 1. \end{cases};$$

$$\begin{cases} \operatorname{arctg} x = t, \\ \arcsin y = t; \end{cases} \Leftrightarrow \begin{cases} x = \operatorname{tg} t, \\ y = \sin t. \end{cases}; 0 < t \leq \frac{\pi}{2} \Rightarrow \sin t = \frac{1}{\sqrt{1 + \operatorname{ctg}^2 t}} \Rightarrow y = \frac{1}{\sqrt{1 + x^2}}.$$

$$2. \operatorname{arctg} x = \arccos y = t;$$

$$\begin{cases} 0 < \operatorname{arctg} x < \frac{\pi}{2}, \\ 0 \leq \arccos y \leq \pi; \end{cases} \Rightarrow 0 < t < \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 < \operatorname{arctg} x < \frac{\pi}{2}, \\ 0 < \arccos y < \pi; \end{cases} \Leftrightarrow \begin{cases} x \in \mathbb{R}, \\ -1 < y < 1. \end{cases};$$

$$\begin{cases} \operatorname{arctg} x = t, \\ \arccos y = t; \end{cases} \Leftrightarrow \begin{cases} x = \operatorname{tg} t, \\ y = \cos t. \end{cases}; 0 < t < \frac{\pi}{2} \Rightarrow \cos t = \frac{\operatorname{ctg} t}{\sqrt{1 + \operatorname{ctg}^2 t}} \Rightarrow y = \frac{x}{\sqrt{1 + x^2}}.$$

$$3. \operatorname{arctg} x = \operatorname{arctg} y = t;$$

$$\begin{cases} 0 < \operatorname{arctg} x < \frac{\pi}{2}, \\ -\frac{\pi}{2} < \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Rightarrow 0 < t < \frac{\pi}{2} \Leftrightarrow \begin{cases} 0 < \operatorname{arctg} x < \frac{\pi}{2}, \\ 0 < \operatorname{arctg} y < \frac{\pi}{2}; \end{cases} \Leftrightarrow \begin{cases} x > 0, \\ y > 0. \end{cases};$$

$$\begin{cases} \operatorname{arctg} x = t, \\ \operatorname{arctg} y = t; \end{cases} \Leftrightarrow \begin{cases} x = \operatorname{tg} t, \\ y = \operatorname{tg} t. \end{cases}; \operatorname{tg} t = \frac{1}{\operatorname{ctg} t} \Rightarrow y = \frac{1}{x}.$$